

FEATURES OF ANTIOXIDANT THERAPY IN PATIENTS WITH CHRONIC
NEPHROTIC GLOMERULONEPHRITIS.

Turaeva Nazira Yuldashevna

Ergashev Abdurashit Khursandovich

Assistants department of Pediatrics-2 of Samarkand State Medical University, Samarkand,
Uzbekistan

Abstract. According to the modern concept, pathological processes are realized on cell membranes, causing a violation of the structural and functional organization, up to complete destruction. The use of glucocorticoid and cytotoxic therapy in combination with heparin does not guarantee success and lead to complications, especially in children. Consequently, the search for drugs that remove the toxic effects of immunosuppressive therapy continues to be an urgent medical and social problem. What was the reason to study the features of antioxidant therapy in order to optimize the pathogenetic treatment of chronic glomerulonephritis in children.

The advantage of therapy with the inclusion of actovegin antioxidant was that parallel to the normalization of malondialdehyde and lysophosphatidylcholine, the other cell membrane fractions increased, apparently due to the ability of aktovegin to reduce the activity of phospholipases. The conducted research will allow to review the tactics of treatment of nephropathy in accordance with the obtained research results, to include in the complex therapy a step-by-step correction of membrane destabilization with Actovegin. In addition, pathogenetic therapy for glomerulonephritis is fraught with complications, ultimately associated with a violation of cellular stability and requires corrective antioxidant therapy.

Key words: glomerulonephritis, nephrotic form, antioxidant therapy.

INTRODUCTION

According to surveys among children, acute glomerulonephritis is 1.1 per cent, chronic at 0.4 per cent. Glomerulonephritis in children tends to increase with the development of chronic kidney failure [1]. High temperatures of up to 41 degrees and above up to 110 days per year, with low relative humidity of up to 20% in Uzbekistan, result in adaptive shifts in the kidneys. Lipid peroxidation activity (LIPID PEROXIDATION) is increased, free radicals concentration is increased as a result of prolonged exposure, antioxidants are depleted [7]. The achievements of modern clinical membranology make it clear that the most important mechanism for regulating the state of cell membranes during adaptation and under stress during the development of various diseases, is the peroxidation of lipids of unsaturated fatty acids and phospholipids flowing along a free radical path. According to the modern concept, pathological processes are realized on the cell membranes, causing structural and functional disruption, up to peroxidation lipid destruction. The volatile oxidation of lipidoformation by glucocorticoid and cytotoxic therapy combined with heparinotheapia do not guarantee success and lead to complications, especially in childhood [2]. Consequently, the search for drugs that relieve the toxic effects of immunosuppressant therapy continues to be a pressing medical and social problem. [3].

The purpose of this work was to determine the effectiveness and indications for the prescription of antioxidant therapy in children with chronic glomerulonephritis nephrotic form.

Research materials and methods. Studies were conducted on 125 patients with chronic glomerulonephritis. The Monitoring Group served 31 practically healthy children of the same age who were not affected by nephropathy. The content of general lipids, cholesterol, concentration of phospholipids by the dencytometer «Byan» by M.H.Turakulov, fraction of phospholipids by E.Stahl method in modification of V.I.Krylov and non-esterified fatty acids, the activity of peroxidation of lipids, determined by the indicator of malonon dialdehyde, the activity of phospholipase A² method. A study of healthy children has been conducted to determine the impact of climate on membrane health. As a result, a higher content of lysophosphatidylcholin, the most toxic fraction of phospholipids, was found in the membrane in summer. Compensatorively, in the summer increases the content of phosphatidylethanolamine, osfatidil ethanolamine, sphingomyelin. In healthy children, lipid peroxidation has been found in small dialdehyde and sphingomyelin (0.55); sphingomyelin and phosphatidylethanolamine (0.6); phosphatidylcholine and lysophosphatidylcholine (0.53). In the active stage of chronic glomerulonephritis in the summer, the highest content of Malon dialdehyde was found as an indicator of high lipid peroxidation activity and was accompanied by an increase in lysophosphatidilcholine content in cell mebranes.

The increase in lipid peroxidation activity in summer was also accompanied by high blood phospholipase activity in patients with a nephrotic form of chronic glomerulonephritis compared to winter. Lipid peroxidation correlated with lysophosphatidylcholine and phosphatidyl ethanolamine (0.82), lysophosphospatidilcholine with phosphatidylcholine (0.74), lysophosphospatidilcholine with phosphatidylserine (0.8). phosphatidylserine with phosphatidylcholine (0.8) is active in patients with glomerulonephritis. In the remission phase, the association of lysophosphathyldilcholine with other phospholipid fractions disappears. In this regard, Aktovegin is included in our complex therapy. The drug is selected as an antioxidant, due to the content of superoxide-dismutase, (72.8 7.1 ml) is characterized by high antioxidant activity compared to other antioxidants. In addition, aktovegin contains macro and trace elements that are part of the enzymes involved in oxidative processes. During the activity of glomerulonephritis intravenously injected 600 - 2000 mg/day in/in drip 10 - 14 days, then transferred to a drage of 200 - 400 mg 2 times per day per os - 1-2 months Dosing of the drug was carried out depending on the age The course was repeated 3-4 times a year if necessary without a break during the year depending on the dynamics of the process.

Research results. Thus, against the background of pronounced clinical manifestations and urinary syndrome in patients with chronic glomeroelecteritis in the nephrotic form increases the intensity of lipid peroxidation. In peroxidation lipidovyse lipid oxidation of such pre-oxidation lipid oxidation also says the identified lipid lipid oxidation bond of the malone dialdehyde and LF during the aggravation period ($h=0.476$; $p < 0.05$). And in remission, his activity decreases. The dynamics of the spectrum of phospholipids with the predominant accumulation of unsaturated phospholipids in lipid bislu indicates the absence of lipid peroxidation in the cytomembrane state in patients with chronic nephrotic glomeroenlytis. Lipid peroxidation correlated with lysophosphatidylcholine ($h=0.465$; $p < 0.01$), negative correlation of total phospholipids and sphingomyelin ($h=0.661$; $p < 0.01$), indicating thatthat when lipid peroxidation activity increases, lysophosphatidyllin and sphingomyelin increase, while total phospholipids decrease. In remission of glomerulonephritis membrane peroxidation lipid does not restore the structure of the membrane of healthy children, and the phospholipids of cell membranes are in a condition that dynamically

determines cellular preservation under peroxidation conditions... In remission, LF,FS,oddnako in parallel increases phosphatidyl ethanolamine to 0.34 0.02, $p < 0.05$ and remains elevated FH(0.34 0.03 mmol/l). Clinical and laboratory remission of patients with chronic glomeronephritis increased the level of small dialdehyde and lysophosphatidylcholine and reduced the number of total phospholipids, which may indicate the activity of the disease. We associate these changes with the fact that as a result of reduced activity, lipid peroxidation during remission produces phospholipid synthesis, while reducing the amount of total phospholipids in plasma,negative correlation of total phospholipids with lysophosphatidylcholine ($h=0.401,p < 0.05$), phosphatidylserine($h=0.491,p < 0.05$), sphingomyelin ($h=-0.481,p < 0.05$) during the remission period, that is, the inflammatory oxidation of the lipid-free defect of the oxidized fractions leads to a decrease in the total phospholipids.

CONCLUSIONS

1. State analysis of lipid peroxidation and spectrum of erythrocyte membrane phospholipids found high lipid peroxidation activity during the exacerbation period with increased lysophosphatidylcholine fraction, reduction of total plasma phospholipids,during the remission-absence of cell membrane stabilization against the background of reduced lipid peroxidation intensity and a decrease in the malone dialdehyde, in patients with chronic nephrotic glomeronephritis.

2. The ratios we've discovered in the spectrum of phospholipids are explained by the change in their oxidation rate, which allows the cell metabolism to be restructured in the membrane in response to various effects, the phospholipid mobility to be lost, significant in the chronology of glomerulonephritis and renal deficiency.

REFERENCES:

1. Batirbekovich, K. R., Sabrieva, V. A., & Alamovich, K. A. (2022). Psychopharmacotherapy of Depressive Disorders in Alcoholism. Eurasian Journal of Humanities and Social Sciences, 8, 19-22.
2. Khayatov, R. B., Velilyaeva, A. S., & Kurbanov, A. A. (2022). OPTIMIZATION OF THERAPY OF ALCOHOL WITHDRAWAL SYNDROME IN PATIENTS WITH SUB-DEPRESSION. Евразийский журнал медицинских и естественных наук, 2(5), 189–192.
3. Khayatov, R. B., Velilyaeva, A. S., & Kurbanov, A. A. (2022). AFFECTIVE DISORDERS AS A WEIGHTENING FACTOR IN ALCOHOL DEPENDENCE THERAPY. Евразийский журнал медицинских и естественных наук, 2(5), 193–196.
4. Telmanovna, X. S., & Batirbekovich, X. R. (2023). Psychopharmacotherapy of Depressive Disorders in Alcoholism. Eurasian Research Bulletin, 16, 179-182.
5. N. Turaeva (2023). CLINICAL-LABORATORY FEATURES OF INTERSTITIAL NEPHRITIS IN CHILDREN WITH PURINE DYSMETABOLISM. Science and innovation, 2 (D12), 135-140. doi: 10.5281/zenodo.10324931
6. N. Turaeva (2023). ANTIOXIDANT THERAPY IN PATIENTS WITH CHRONIC NEPHROTIC GLOMERULONEPHRITIS. Science and innovation, 2 (D12), 131-134. doi: 10.5281/zenodo.10324779
7. Меликова Дилшодахон Уктам Кизи, Ахмеджанова Наргиза Исмаиловна, Тураева Назира Юлдашевна, Юлдашев Ботир Ахматович, & Абдурасулов Фозил Пардаевич (2020). Клинические особенности течения хронического пиелонефрита у детей на фоне анемического синдрома. Достижения науки и образования, (1 (55)), 66-69.

8. Очилов, У., Кубаев, Р., & Хаятов, Р. (2016). Психические расстройства при употреблении психоактивных веществ с вич-инфекцией. Журнал проблемы биологии и медицины, (2 (87)), 184–186.
9. Раджабов Хикмат Тошевич, Хаятов Рустам Батырбекович, & Велиляева Алие Сабриевна (2020). КЛИНИКО-ПСИХОЛОГИЧЕСКИЕ И НЕЙРОФИЗИОЛОГИЧЕСКИЕ ОСОБЕННОСТИ НЕПСИХОТИЧЕСКИХ ПСИХИЧЕСКИХ РАССТРОЙСТВ У ЛИЦ ПОЖИЛОГО ВОЗРАСТА. Вестник науки и образования, (23-3 (101)), 75-78.
10. Раджабов Х.Т., Тургунбаев А.У., Кубаев Р.М., & Хаятов Р.Б. (2021). ДЕПРЕССИЯ И ТРЕВОГА У БОЛЬНЫХ АЛКОГОЛИЗМОМ, ОСЛОЖНЕННЫМ НИКОТИНОВОЙ ЗАВИСИМОСТЬЮ. Вестник науки и образования, (17-2 (120)), 134-137.
11. Тураева Назира Юлдашевна (2020). Клинико-лабораторные особенности течения дисметаболической нефропатии у детей с нарушением пуринового обмена. Достижения науки и образования, (5 (59)), 86-88.
12. Тураева, Н., & Абдукадырова, Н. (2022). Оптимизация терапии хронического гломерулонефрита у детей. Журнал вестник врача, 1(2), 118–120.
13. Хаятов Рустам Батырбекович, Велиляева Алие Сабриевна, Тураев Бобир Темирпулатови, & Тураев Толиб Махмуджонович (2019). Аффективные расстройства у больных алкогольной зависимостью как фактор риска развития суицидального поведения. Достижения науки и образования, (11 (52)), 96-98.
14. Хаятов Рустам Батырбекович, & Велиляева Али Сабриевна (2020). Особенности развития и течения аффективных расстройств при сахарном диабете. Достижения науки и образования, (5 (59)), 62-64.
15. Хаятов Рустам Батырбекович, Велиляева Алие Сабриевна, & Абдуразакова Робия Шералиевна (2020). Особенности возникновения и течения психоорганических расстройств при сахарном диабете. Достижения науки и образования, (7 (61)), 31-33.
16. Хаятов, Р., & Велиляева, А. (2022). Влияние тревожно-депрессивных расстройств на тяжесть течения и качество жизни у больных сахарным диабетом 2 типа. Журнал вестник врача, 1(4), 99–102. <https://doi.org/10.38095/2181-466X-2020974-98-101>
17. Хаятов, Р., Велиляева, А., Мардиев О., & Курбанов, А. (2022). Особенности коморбидного течения тревожно-депрессивных расстройств и личностных изменений при сахарном диабете 2 типа. Журнал вестник врача, 1(1), 104–108. <https://doi.org/10.38095/2181-466X-2021981-103-107>
18. Каршиев Зиядулло Хазратович, Хаятов Рустам Батырбекович, Шерматов Озод Норбекович, Раджабов Хикмат Тошевич, & Рузиева Диана Джамаловна (2021). АФФЕКТИВНЫЕ РАССТРОЙСТВА КАК ОТЯГОЩАЮЩИЙ ФАКТОР В ТЕРАПИИ АЛКОГОЛЬНОЙ ЗАВИСИМОСТИ. Вестник науки и образования, (5-2 (108)), 21-24.
19. Хаятов, Р., Велиляева, А., Мардиев О., & Курбанов, А. (2022). Особенности коморбидного течения тревожно-депрессивных расстройств и личностных изменений при сахарном диабете 2 типа. Журнал вестник врача, 1(1), 104–108. <https://doi.org/10.38095/2181-466X-2021981-103-107>
20. Очилов, У., Кубаев, Р., & Хаятов, Р. (2016). Психические расстройства при употреблении психоактивных веществ с вич-инфекцией. Журнал проблемы биологии и медицины, (2 (87)), 184–186. извлечено от https://inlibrary.uz/index.php/problems_biology/article/view/3568

21. Раджабов Х.Т., Тургунбаев А.У., Кубаев Р.М., & Хаятов Р.Б. (2021). ДЕПРЕССИЯ И ТРЕВОГА У БОЛЬНЫХ АЛКОГОЛИЗМОМ, ОСЛОЖНЕННЫМ НИКОТИНОВОЙ ЗАВИСИМОСТЬЮ. Вестник науки и образования, (17-2 (120)), 134-137.
22. Хаятов, Р., & Абдуразакова, Р. (2023). Аффективные расстройства, как отягощающий фактор в терапии алкогольной зависимости. Журнал биомедицины и практики, 1(3/1), 396–399. извлечено от <https://inlibrary.uz/index.php/biomedicine/article/view/18375>
23. Мардиев, О. ., & Хаятов, Р. (2023). КОМОРБИДНОСТЬ ДЕПРЕССИВНЫХ РАССТРОЙСТВ И САХАРНОГО ДИАБЕТА 2-го ТИПА НА КАЧЕСТВО ЖИЗНИ БОЛЬНЫХ ПОЖИЛОГО ВОЗРАСТА. *Евразийский журнал медицинских и естественных наук*, 3(8), 19–24. извлечено от <https://in-academy.uz/index.php/EJMNS/article/view/19616>
24. Николаев Егор Евгеньевич, Орлов Федор Витальевич, Николаев Евгений Львович, Велиляева Алие Сабриевна, & Хаятов Рустам Батырбекович (2023). ФЕНОМЕН СИМУЛИРОВАНИЯ СУДОРОЖНЫХ ПРИСТУПОВ ПРИ ЭПИЛЕПСИИ: КЛИНИЧЕСКИЙ СЛУЧАЙ. *Acta Medica Eurasica*, (3), 102-115. doi: 10.47026/2413-4864-2023-3-102-115
25. Criteria For Rehabilitation Of Patients With Consequences Cranio-Brain Injury. (2022). *Journal of Pharmaceutical Negative Results*, 8188-8194. <https://doi.org/10.47750/pnr.2022.13.S09.958>
26. Хаятов, Р. Б., & Велиляева, А. С. НЕЙРОПСИХОЛОГИЧЕСКОЕ ИССЛЕДОВАНИЕ БОЛЬНЫХ С РАССЕЯННЫМ СКЛЕРОЗОМ. ДОКТОР АХБОРОТНОМАСИ ВЕСТНИК ВРАЧА DOCTOR'S HERALD, 44.
27. Мардиев Отабек Аслиддинович, Кубаев Рустам Мурадиллаевич, Хаятов Рустам Батырбекович, & Рузиева Диана Джамаловна (2021). ТЕЧЕНИЕ ТРЕВОЖНЫХ РАССТРОЙСТВ У БОЛЬНЫХ САХАРНЫМ ДИАБЕТОМ 2 ТИПА. Вестник науки и образования, (2-2 (105)), 72-75.
28. Велиляева, А., Бердиева, Н., Очилов, У., & Хаятов, Р. (2015). Психические нарушения при эпилепсии. *Журнал проблемы биологии и медицины*, (2 (83)), 168-171.
29. Депрессия и тревога у больных алкоголизмом, осложненным никотиновой зависимостью / Х. Т. Раджабов, А. У. Тургунбаев, Р. М. Кубаев, Р. Б. Хаятов // Вестник науки и образования. – 2021. – № 17-2(120). – С. 134-137. – EDN GSEZVQ.
30. Turaeva, N. Y., & Yuldashev, V. A. (2018). KLINIKO-LABORATORNYE POKAZATELI INTERSTITsIAL'NOY PATOLOGII POCHEK U DETEY V STRUKTURE DISMETABOLIChESKIKh NEFROPATIIY. *Молодежный инновационный вестник*, 7(S1), 99-100.
31. Тураева, Н. Ю. Клинико-лабораторные показатели интерстициальной патологии почек у детей в структуре дисметаболических нефропатий / Н. Ю. Тураева, Б. А. Юлдашев // Молодежный инновационный вестник. – 2018. – Т. 7, № S1. – С. 99-100. – EDN YXJTТА.