

EPIDEMIOLOGY AND COMPLICATIONS OF COVID-19

Ibadullayeva Sevara Salibayevna, Yoqubov Qahhor Yodgor o'g'li, Razzoqberganova Dinora Odilbekovna

Urgench branch of Tashkent Medical Academy, Urgench, Uzbekistan

Annotation: The liver is a vital organ. The liver has many functions in the human body. Participates in the reactions of the immune system. In addition, it determines the overall level of resistance to infections and is responsible for the production of antibodies to fight viruses and bacteria. Liver diseases are the most common in the world. Especially in the context of the COVID-19 pandemic. World Health Organization February 11, 2020.

Keywords: coronavirus, COVID-19, chronic hepatitis, clinic, mutation, lymphocytes, immunity.

Introduction: News by the end of 2019 In the People's Republic of China, an unknown pneumonia has attracted universal attention. Pneumonia caused by the new coronavirus SARS-CoV-2 severe acute respiratory syndrome coronavirus is a coronavirus of severe acute respiratory syndrome. On February 11, 2020, the World Health Organization officially assigned the name SARS-CoV-2 infection, COVID-19 "CoronaVirus Disease 2019" — a disease caused by the new coronavirus 2019). Pneumonia is the leading clinical form of COVID-19.

Nevertheless, as the COVID-19 pandemic spread and the analysis of clinical data, patients began to identify symptoms that were not characteristic of "abnormal" pneumonia. Neurological manifestations, skin changes, eye lesions, etc. are described. The extrapulmonary presence of SARS-CoV-2 was also found in cholangiocytes. Among the possible factors of liver damage, virus-induced effects, systemic inflammation ("cytokine storm"), hypoxia, hypovolemia, hypotension in shock, drug hepatotoxicity, etc. are considered. In 14-53% of patients with COVID-19, changes in biochemical parameters may be registered, which, as a rule, do not require medical correction. Acute hepatitis is very rare. However, special attention should be paid to patients with COVID-19 belonging to the risk group — those who have undergone liver transplantation, receiving immunosuppressants, as well as in cases of decompensation of cirrhosis, the development of acute liver failure against the background of chronic, hepatocellular carcinoma, during antiviral therapy. There is a need for constant exchange and open access to scientific data, new technologies, and updated guidelines.

Coronaviruses are widespread in nature and are the causes of various colds (up to 25%). Most of them cause a viral infection that does not cause serious harm to health, but some, such as SARS-CoV severe acute respiratory syndrome coronavirus — severe acute respiratory syndrome coronavirus and MERS-CoV (Middle East respiratory syndrome coronavirus-Middle East respiratory syndrome coronavirus) lead to the development of severe respiratory syndrome with high mortality. In nature, many species of bats serve as the natural host for coronaviruses.

Evolving due to mutations and preadaptation processes, they periodically cause epidemics in human populations. Thus, the outbreak of an unknown pneumonia in late December 2019 in China caused the development of a public health emergency, which subsequently led to a pandemic caused

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-3

by the new coronavirus SARS-CoV-2 (severe acute respiratory syndrome coronavirus - severe acute respiratory syndrome coronavirus). World Health Organization (WHO) February 11, 2020 It has assigned the official name of SARS-CoV-2 infection, COVID-19 ("CoronaVirus Disease 2019" is a disease caused by the new coronavirus 2019). The mortality rate from this infection is 0.5-3%. SARS-CoV-2 and possible mechanisms of COVID-19 pathogenesis. The new coronavirus is a single-stranded RNA-containing virus belonging to the family Coronaviridae, genus Betacoronavirus. SARS-CoV-2 is a zoonotic virus, which follows from phylogenetic analysis, which showed the closest relationship with the isolate of the SARS-like bat coronavirus BM48-31/BGR/2008 (identity — 96%).

Apparently, bats are the reservoir of SARS-CoV-2, and other small mammals, in particular, pangolins, are intermediate hosts, possibly infecting the "patient zero". In addition, the phylogenetic analysis of SARS-CoV-2 obtained data indicating 88% sequence identity with SARS-CoV and about 50% with MERS-CoV. The structure of coronaviruses of respiratory syndromes is very similar. Among the structural proteins of SARS-CoV-2, S-proteins or "protein spikes" (from the English Spike — spike), membrane protein, protein and nucleocapsid are distinguished. Protein S plays an important role in the attachment, fusion and penetration of the virus into cells, which allows it to be considered as a possible target for the production of antibodies and vaccines. The pathogenesis of the new coronavirus infection has not been sufficiently studied. A key virulence factor is the interaction of the receptor-binding domain (receptor—binding domain - RBD) of protein S located on the outer membrane of SARS-CoV-2 with the receptor of angiotensin-converting enzyme 2 (angiotensin—converting enzyme 2 receptors - ACE2) activated by transmembrane serine proteases (TMPRSS2 - Transmembrane protease, serine 2) of humans. ACE2 is expressed in a surfactant secreted by type II alveolocytes from blood plasma components.

Surfactant is a surfactantly active monomolecular film, which is located at the interface of the air-liquid phases in the alveoli, alveolar passages and respiratory bronchioles of the 1st—3rd order and prevents the collapse (adhesion) of the walls of the alveoli during breathing. The expression of ACE2 protects against lung damage, but it decreases due to its binding to the SARS-CoV spike protein, which increases the risk of infection. At the same time, it was shown in the experiment that increased expression of ACE2 does not exclude the possibility of increased binding to SARS-CoV. Up to three viruses can attach to a single target. ACE2 and TMPRSS2 are unevenly distributed among patients of European and Asian origin, which can also affect the intensity of infection.

It has been suggested that non-structural SARS-CoV proteins are capable of modifying the structure of hemoglobin in the erythrocyte, which leads to a violation of oxygen transport, causes iron dissociation, porphyrin formation, and an increase in ferritin. Such exposure can lead to increased inflammatory processes in the lungs, oxidative stress, hypoxemia, hypoxia, the development of symptoms of acute respiratory distress syndrome ARDS and multiple organ oxygen deficiency. However, the basis for this hypothesis was the creation of a biotransformation model without conducting experimental and clinical studies of SARS-CoV-2 tropens to goblet cells contained in the mucous membrane of the respiratory tract, intestine, conjunctiva of the eyes, ducts of the pancreas and parotid salivary glands. Active replication of the virus significantly reduces the protective functions of goblet cells (mucus formation), which also contributes to the penetration of the virus into the human body.

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-3

In response to the spread of coronavirus, the development of a hyperimmune reaction is observed — the so-called "cytokine storm", characterized by the synthesis of a significant (abnormal) amount of proinflammatory interleukins (IL-1 β , IL-6, tumor necrosis factor, etc.) and chemokines while reducing the content of T-lymphocytes in the blood. In addition, SARS-CoV-2, infecting the endothelium of blood vessels, interacts with ACE2 located there and leads to the development of endothelial dysfunction, hyper permeability, impaired microcirculation, the development of vascular thrombophilia and thrombosis. The progression of COVID-19 is determined by diffuse alveolar damage with the formation of hyaline membranes, the development of pulmonary edema. The histological picture of the lungs during autopsy is characterized by the organization of alveolar exudates and interstitial fibrosis, the formation of hyaline membranes, the presence of interstitial mononuclear inflammatory infiltrates, numerous microthrombs of fibrin, pronounced edema, hyperplasia and focal desquamation of type II alveolocytes, a significant content of macrophages with viral inclusions in the alveolar exudate. Hemorrhage, necrosis, and hemorrhagic infarction are observed in more affected areas. The clinical picture and diagnosis of COVID-19 infection with SARS-CoV-2 in humans occurs in the last days of the incubation period and as much as possible in the first three days from the onset of the disease.

In the vast majority, infection occurs due to contact with a COVID-19 patient in cases of clinically manifested disease (up to 75-85% — in contact with infected relatives from a family environment). It should be emphasized that the release of the virus, as a rule, lasts up to 12 days in mild/moderate cases and more than 14 days in severe cases. However, in patients who have recovered from COVID-19, the RNA of the new coronavirus may be positive even after the disappearance of clinical symptoms. A patient with COVID-19 can infect 3-5 people around him, and a person with the flu can infect only 1-2 people.

Epidemiological data indicate that patients with cardiovascular diseases, hypertension, diabetes mellitus, and malignant tumors are the most susceptible to SARS-CoV-2.

The incubation period is 2-14 days, the average period is 5-6 days. SARS - CoV-2 is transmitted by airborne droplets during coughing, sneezing, talking, airborne dust with dust particles in the air, contact through handshakes, household items and fecal-oral routes.

The identified adverse signs require monitoring of liver function and electrocardiography, especially among risk groups — patients with CKD and myocardial repolarization disorders. The scale in the use of experimental treatments for COVID-19 is unprecedented. However, evidence of their effectiveness has yet to be established. In this regard, the issue of drug interaction remains very important for clinical practice. On the website of the University of Liverpool (UK) (www.covid19-druginteractions.org) the main experimental drugs that are currently used in COVID-19 therapy are presented, indicating their mechanisms of action; an assessment of joint use with other drugs is given, taking into account the risks and benefits, duration of use, patient's condition, and medication for previously established diseases.

Conclusion:

There are over 3.5 million cases of COVID-19 worldwide, which has killed more than 250,000 people. Unfortunately, there are currently no effective specific treatments for COVID-19. Numerous clinical randomized trials of various drugs are being conducted. So far, there is no evidence that

patients who have recovered from COVID-19 are protected from re-infection. It is necessary to monitor individuals with antibodies against SARS-CoV-2 in comparison with those without them, with an assessment of the frequency of infection with SARS-CoV-2 and the development of COVID-19 over a long period (at least one year). However, the first experimental experience of using plasma containing class G immunoglobulin antibodies in patients who have undergone COVID-19 has shown encouraging results. Recovered patients from COVID-19 and people with an asymptomatic course who secrete the virus with faeces can be considered as a possible source of infection. In addition, since SARS-CoV-2 RNA has been detected in wastewater samples, the question of the viability of the virus in an environment through which a fecal-oral transmission pathway can also be implemented remains unresolved.

List of literature.

1. Masharipov, S., Sadullaev, S. E., & Sh, M. D. (2023). THE COURSE OF CORONAVIRUS AGAINST THE BACKGROUND OF CHRONIC HEPATITIS. *Scientific Impulse*, 2(15), 65-70.
2. Masharipova Sh.S, Masharipov S, Sadullaev S.E, & Matyakubova D.Sh. (2023). THE COURSE OF CORONAVIRUS AGAINST THE BACKGROUND OF CHRONIC HEPATITIS. *Scientific Impulse*, 2(15), 65–70. Retrieved from
3. Sadullayev S. E. et al. THE COURSE OF NOSOCOMIAL PNEUMONIA IN PATIENTS ON LONG-TERM ARTIFICIAL LUNG VENTILATION //O'ZBEKISTONDA FANLARARO INNOVATSIYALAR VA ILMIY TADQIQOTLAR JURNALI. – 2024. – Т. 2. – №. 26. – С. 80-84.
4. Машарипова Шохиста Сабировна, Артиков Икром Ахмеджанович, Матякубова Ойша Уриновна РАСТРОЙСТВА ПСИХИКИ У БОЛЬНЫХ ДЕСТРУКТИВНЫМИ ФОРМАМИ ТУБЕРКУЛЕЗА В УСЛОВИЯХ ПАНДЕМИИ COVID-19 // JCRR. 2021. №3.
URL: <https://cyberleninka.ru/article/n/rastroystva-psihipiki-u-bolnyh-destruktivnymi-formami-tuberkuleza-v-usloviyah-pandemii-covid-19>
5. Машарипова Шохиста Сабировна МОРФОЛОГИЧЕСКОЕ СТРОЕНИЕ ЛЕГОЧНЫХ АРТЕРИЙ ПОД ВЛИЯНИЕМ САХАРНОГО ДИАБЕТА // JCRR. 2022. №1.
URL: <https://cyberleninka.ru/article/n/morfologicheskoe-stroenie-legochnyh-arteriy-pod-vliyaniem-saharnogo-diabeta>
6. Туйчиев Л. Н. и др. A study of the factors affecting the effectiveness of COVID-19 rehabilitation. – 2023.
7. Машарипова Ш. С. О ‘РКА АРТЕРИЯЛАРИНИНГ QANDLI ДИАБЕТ ТА’СИРИДА MORFOLOGIK TUZILISHI //Журнал кардиореспираторных исследований. – 2022. – Т. 3. – №. 1.
8. Yusupov Sh.Sh, Ibrakhimova H.R, & Masharipova Sh.S. (2023). IMMUNOLOGICAL CHARACTERISTICS OF PATIENTS WHOSE BODY IS INFECTED WITH CATTLE SOLITAIRE. *Новости образования: исследование в XXI веке*, 2(15), 120–124. извлечено от <http://nauchniyimpuls.ru/index.php/noiv/article/view/12430>

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-3

9. Ibrakhimova H.R, Masharipova Sh.S, & Artikov I.A. (2023). CHANGES IN THE IMMUNE STATUS OF PATIENTS WITH PARASITIC DISEASE. *Новости образования: исследование в XXI веке*, 2(15), 103–108. извлечено от <http://nauchniyimpuls.ru/index.php/noiv/article/view/12427>
10. Masharipova Sh.S, Masharipov S, & Matyakubova D.Sh. (2023). UDC:616.36-005:75-642 TUBERCULOSIS AND ITS COURSE IN PATIENTS WITH HEPATITIS B. *Scientific Impulse*, 2(14), 95–99. Retrieved from <http://nauchniyimpuls.ru/index.php/ni/article/view/12215>
11. Masharipova Sh.S, Ibrakhimova H.R, & Nurlayev R.R. (2023). A METHOD FOR OBTAINING PRECIPITATING SERUMS FOR THE DETECTION OF HUMAN SEMINAL FLUID USED IN THE STUDY OF PHYSICAL EVIDENCE IN FORENSIC BIOLOGICAL LABORATORIES. *World Bulletin of Management and Law*, 19, 42-44. Retrieved from <https://scholarexpress.net/index.php/wbml/article/view/2119>
12. IMMUNE STATUS IN PATIENTS WITH PARASITIC DISEASES IN KHOREZM REGION. (2025). *Multidisciplinary Journal of Science and Technology*, 5(1), 514-517. <http://www.mjstjournal.com/index.php/mjst/article/view/2518>
13. Машарипов С.М., Юсупов Ш.Р., Машарипова Ш.С., Матякубова О.У. КЛИНИЧЕСКОЕ ТЕЧЕНИЕ ТУБЕРКУЛЕЗА У БОЛЬНЫХ ГЕПАТИТОМ В / Вестник ТМА.uz. 2023 г. № 3 (2) – стр. 155-157.
14. Аскарлова Р.И., Юсупов Ш.Р., Машарипова Ш.С., Машарипова Х.К. Эпидемиология легочного туберкулеза/EUROPEAN RESEARCH: INNOVATION IN SCIENCE, EDUCATION AND TECHNOLOGY LVII International correspondence scientific and practical conference. 2019/Издательство: PROBLEMS OF SCIENCE / United Kingdom, 06–07 ноября 2019 год, стр. 96-100
15. S., M. S., E., S. S. , J., I. S. , I., Q. N. , & B., B. Y. . (2024). THE SPREAD OF CHICKENPOX AMONG THE POPULATION. *Research Journal of Trauma and Disability Studies*, 3(3), 48–54. Retrieved from <http://journals.academiczone.net/index.php/rjtds/article/view/2302>
16. Masharipova Sh.S, Ibadullayeva S.S, Yakubov K.Y, & Artikov I.A. (2024). FEATURES OF INFECTIOUS MONONUCLEOSIS IN CHILDREN. *Scientific Impulse*, 2(16), 1165–1171. Retrieved from <http://nauchniyimpuls.ru/index.php/ni/article/view/14208>
17. Masharipova Sh.S, Ibadullayeva S.S, Yakubov K.Y and Artikov I.A 2024. FEATURES OF INFECTIOUS MONONUCLEOSIS IN CHILDREN. *Scientific Impulse*. 2, 16 (Jan. 2024), 1165–1171.
18. Садуллаев, С. Е., Машарипова, Ш. С., & Машарипов, С. (2023). КЛИНИКО-ЛАБОРАТОРНЫЕ ОСОБЕННОСТИ ТЕЧЕНИЯ ПНЕВМОНИИ, АССОЦИИРОВАННОЙ С COVID-19, У ДЕТЕЙ РАННЕГО ВОЗРАСТА. *Международный журнал образования, социальных и гуманитарных наук*. Finland Academic Research Science Publishers, 11(9), 851-856.

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-3

19. Сабировна, Ш., & Машарипова, А. И. А. Садуллаев Сирож Эрназарович, и Абдуллаева Дилфуза Кадамовна. 2022. «ТЕЧЕНИЕ КОРОНАВИРУСНОЙ ИНФЕКЦИИ НА ФОНЕ ГЕПАТИТОВ». *Новости образования: исследование в XXI веке* 1 (5): 573-77. ТЕЧЕНИЕ КОРОНАВИРУСНОЙ ИНФЕКЦИИ НА ФОНЕ ГЕПАТИТОВ». *Новости образования*.
20. Masharipova Sh.S, Masharipov S, & Matyakubova D.Sh. (2023). UDC:616.36-005:75-642 TUBERCULOSIS AND ITS COURSE IN PATIENTS WITH HEPATITIS B. *Scientific Impulse*, 2(14), 95–99. Retrieved from <https://nauchniyimpuls.ru/index.php/ni/article/view/12215>
21. Masharipova Sh.S, Masharipov S, & Matyakubova D.Sh. (2023). UDC:616.36-005:75-642 TUBERCULOSIS AND ITS COURSE IN PATIENTS WITH HEPATITIS B. *Scientific Impulse*, 2(14), 95–99. Retrieved from <http://nauchniyimpuls.ru/index.php/ni/article/view/12215>
22. Masharipova Shokhista Sabirovna, & Masharipov Sobir. (2023). UDC: 619:616.995.132.6 IMMUNE STATUS OF ADULTS AND CHILDREN WITH AN ALLERGIC BACKGROUND DIAGNOSED WITH ENTEROBIOSIS. *Новости образования: исследование в XXI веке*, 2(14), 24–28. извлечено от <https://nauchniyimpuls.ru/index.php/noiv/article/view/11911>
23. CHARACTERISTICS OF PATHOMORPHOLOGICAL CHANGES IN LYMPHOCYTIC LEUKOSIS IN CHILDREN. (2023). *Western European Journal of Medicine and Medical Science*, 1(4), 21-26. <https://westerneuropeanstudies.com/index.php/3/article/view/122>
24. RESULTS OF STUDIES ON THE LEVEL OF POPULATION KNOWLEDGE ABOUT PARASITIC DISEASES AND ITS PREVENTION. (2023). *Western European Journal of Medicine and Medical Science*, 1(4), 15-20. <https://westerneuropeanstudies.com/index.php/3/article/view/121>
25. THE STRUCTURE OF COMORBID PATHOLOGY IN CHILDREN WITH COVID-19. (2024). *CONFERENCE ON THE ROLE AND IMPORTANCE OF SCIENCE IN THE MODERN WORLD*, 1(2), 27-28.
26. Masharipov S., Sadullaev S. E., Sh M. D. THE COURSE OF CORONAVIRUS AGAINST THE BACKGROUND OF CHRONIC HEPATITIS //Научный импульс. – 2023. – С. 78.
27. Masharipova Sh.S., Ibrakhimova H.R, Sadullaev S.E., & Nurllayev R.R. (2023). SPREAD OF MYOCARDIAL INFARCTION AMONG THE POPULATION OF THE KHOREZM REGION. *IMRAS*, 6(7), 328–332. Retrieved from <https://journal.imras.org/index.php/sps/article/view/523>
28. Ibraximova H. R., Sadullayev S. E. AHOLI ORASIDA O ‘TKIR ICHAK KASALLIKLARINING TARQALISHI //Новости образования: исследование в XXI веке. – 2023. – Т. 2. – №. 15. – С. 115-119.
29. Ibrakhimova, H. R., Matyakubova, O. U., Sadullaev, S. E., & Abdullayeva, D. K. (2023). *HELMINTISES IN CHILDREN AMONG THE POPULATION IN UZBEKISTAN. IMRAS*, 6 (7), 323–327.

THE MULTIDISCIPLINARY JOURNAL OF SCIENCE AND TECHNOLOGY

VOLUME-5, ISSUE-3

30. Artikov I.A, Sadullaev S.E, Ibrakhimova H.R, & Abdullayeva D.K. (2023). RELEVANCE OF VIRAL HEPATITIS EPIDEMIOLOGY. *IMRAS*, 6(7), 316–322. Retrieved from <https://journal.imras.org/index.php/sps/article/view/520>

31. Artikov, I. A., Sadullaev, S. E., Ibrakhimova, H. R., & Abdullayeva, D. K. (2023). *RELEVANCE OF VIRAL HEPATITIS EPIDEMIOLOGY. IMRAS*, 6 (7), 316–322.

32. Tychiev, L. N., Khudaykulova, G. K., Eraliev, U. E., Djuraeva, N. K., & Sadullaev, S. E. (2023). A STUDY OF THE FACTORS AFFECTING THE EFFECTIVENESS OF COVID-19 REHABILITATION.

